

INTERESTING FACTS ABOUT SUICIDE

Presentation for Interns at Four County- Logansport

Suicide is the ___ leading cause of death in the U.S.	Suicide risk can be highest when a person has just been discharged from the hospital. T or F?	People who seek treatment are more/less likely to complete suicide?	Which group attempts suicide more often—men or women?	Suicide is the ___ leading cause of death for youths between the ages of 15 and 24.
What are two risk factors for attempt suicide in adults?	The lowest suicide rate for all ages combined is ___ women.	The gender ratio for suicide (M:W) is ___.	Substance use increases the likelihood of suicide risk. T or F	More than 90% of people who kill themselves have some diagnosable mental or substance abuse disorder. T or F?
What is the most common method of suicide? For men? For women?	College students are more/less at risk for suicide than non-students the same age?	People with personality disorders are ___ times as likely to die by suicide than those without.	LGBTIQ issues and gender identity concerns are risk factors for suicide. T or F	People who die by suicide are often suffering from undiagnosed, undertreated, or untreated depression. T/F?
What percentage of people diagnosed w/ schizophrenia die by suicide?	What are two warning signs of suicide?	APA has developed practice guidelines for assessing and treating patients with suicide behaviors. T or F	In depth discussion of suicidal ideation increases the likelihood of follow through by the client. T or F?	Creating a no-suicide contract with a client decreases the likelihood of an attempt or suicide. T or F?

09/2015